



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Substituting this value of x in the last two fractions,

$$\frac{dz}{a} = \frac{dy}{1+cy}, \text{ or } z = \frac{a}{c} \log(1+cy) + c'.$$

Replacing c by $\frac{1}{x} - \frac{1}{y}$, $z = \frac{axy}{y-x} \log \frac{y}{x} + \phi\left(\frac{1}{x} - \frac{1}{y}\right).$

Also solved similarly by J. Scheffer, and G. B. M. Zerr.

293. Proposed by V. M. SPUNAR, M. and E. E., East Pittsburg, Pa.

Find the length of the integral curve of the differential equation
 $(y^2 x^{\frac{1}{3}} + 2)dx - x^{\frac{1}{3}} dy = 0$ between $x_1 = 1$ and $x_2 = 8$.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Let $y = -1/z$, $x^{\frac{1}{3}} = v$, then the equation becomes $dz + 6z^3 dv + 3v^{-4} dv = 0$.

Now, let $z = \frac{1}{6v} + \frac{u}{v^2}$.

Then $v^2 du + 6u^2 dv + 3dv = 0$, or $\frac{du}{3 + 6u^2} = -\frac{dv}{v^2}$.

$$\therefore \frac{1}{3\sqrt{2}} \tan^{-1}(u\sqrt{2}) = \frac{1}{v} + a = a + x^{\frac{1}{3}} \dots (1).$$

$$\therefore u = \frac{1}{\sqrt{2}} \tan[3\sqrt{2}(a + x^{\frac{1}{3}})] \dots (2).$$

$$\therefore y = -\frac{6\sqrt{2}}{6x^{\frac{1}{3}} \tan[3\sqrt{2}(a + x^{\frac{1}{3}})] + x^{\frac{1}{3}}\sqrt{2}} \text{ is the equation.}$$

$$\text{From (1), } u = \frac{1}{\sqrt{2}} \tan\left[\frac{3\sqrt{2}(av+1)}{v}\right].$$

$$S = \int \sqrt{[1 + (du/dv)^2]} dv = \int_{\frac{1}{2}}^1 \frac{1}{v^2} \sqrt{v^4 + 9 \sec^2 \left[\frac{3\sqrt{2}(av+1)}{v} \right]} dv.$$

294. Proposed by C. N. SCHMALL, New York City.

Examine the function, $f(x) = \frac{(x-1)(x-2)}{(x-3)}$ and determine why its *minimum* value is *greater* than its maximum.

Solution by PROF. F. L. GRIFFIN, Ph. D., Williams College.

The derivative, $f'(x) = \frac{x^2 - 6x + 7}{(x-3)^2}$, vanishes for $x = 3 \pm \sqrt{2}$, changing its sign from + to - for the smaller of these values and from - to + at the larger. Thus the function decreases from its maximum value $f(3 - \sqrt{2})$ to its minimum $f(3 + \sqrt{2})$, *except that it passes through* ∞ at $x = 3$, and thus it is possible for the minimum to exceed the maximum. The same fact holds for the more general function of the same type: $F(x) = \frac{(x-a)(x-b)}{x-c}$, if $c > a$ and $c > b$; but the maximum of $F(x)$ exceeds the minimum if $c < a$ and $c < b$. Thus the property proposed for explanation depends not so much on the mere numerical values, nor even on the fact of an infinite discontinuity, as upon the order of the zeros of the numerator and denominator.

Also solved by S. G. Barton, J. E. Sanders, V. M. Spunar, and J. Scheffer.

MECHANICS.

245. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

A body moves with constant speed in the circumference of an ellipse. Find the rate of approach (1) to the center, (2) to one of the foci, for any point in the ellipse.

Solution by PROF. F. L. GRIFFIN, Ph. D., Williams College.

Differentiating $b^2x^2 + a^2y^2 = a^2b^2$ with respect to the time, we have $\frac{dx}{dt}/a^2y = -\frac{dy}{dt}/b^2x = -\lambda$ (say), whence if k denote the constant speed, $k^2 = \left(\frac{ds}{dt}\right)^2 = \left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 = \lambda^2 [a^4y^2 + b^4x^2]$, or $\lambda = k/\sqrt{[a^4y^2 + b^4x^2]}$. The negative sign should be taken if the motion is clockwise.

Hence, $\frac{dx}{dt} = -ka^2y/\sqrt{[a^4y^2 + b^4x^2]}$ and $\frac{dy}{dt} = kb^2x/\sqrt{[a^4y^2 + b^4x^2]}$.

(I) Now the radius vector from the center to (x, y) is given by $r^2 = x^2 + y^2$; whence the rate of approach to the center is

$$-\frac{dr}{dt} = -\left(x\frac{dx}{dt} + y\frac{dy}{dt}\right)/r = -\frac{kxy(b^2 - a^2)}{r\sqrt{[a^4y^2 + b^4x^2]}} = \frac{ka^2e^2xy}{\sqrt{[(a^4y^2 + b^4x^2)(x^2 + y^2)]}}.$$

(II) The radius vector from the left-hand focus $(-ae, 0)$ to (x, y) is given by: $r'^2 = (x + ae)^2 + y^2$; whence

$$-\frac{dr'}{dt} = -[(x + ae)\frac{dx}{dt} + y\frac{dy}{dt}]/r' = \frac{ka^2ey(ex + a)}{\sqrt{[a^4y^2 + b^4x^2](x^2 + y^2)}}.$$